arXiv:1207.1846v1 [hep-ph] 8 Jul 2012

IISER-K/HEP/06/12

New limits on top squark NLSP from LHC 4.7 fb^{-1} data

Arghya Choudhury[1] and Amitava Datta[2]

Indian Institute of Science Education and Research - Kolkata,
Mohanpur Campus, PO: BCKV Campus Main Office,
Nadia, West Bengal - 741252, India.

Abstract

Using the ATLAS 4.7 fb^{-1} data on new physics search in the jets + $\not{E}_T$ channel, we obtain new limits on the lighter top squark ($\tilde{t}_1$) mass assuming that it is the next to lightest supersymmetric particle (NLSP). If the decay $\widetilde{t}_1 \to c\widetilde{\chi}_1^0$ dominates and the production of dark matter relic density is due to NLSP - LSP co-annihilation then the lower limit on $m_{\widetilde{t}_1}$ is 240 GeV. The limit changes to 200 GeV if the decay $\widetilde{t}_1 \to bW\widetilde{\chi}_1^0$ dominates. Combining these results it follows that $\widetilde{t}_1$ NLSP induced baryogenesis is constrained more tightly.

---

[1] arghyac@iiserkol.ac.in

[2] adatta@iiserkol.ac.in

Models with Supersymmetry (SUSY) [1] are very well motivated extensions of the Standard model (SM) of particle physics. The LHC experiments [2, 3] with 4.7 fb^{-1} data have severely constrained the masses of the strongly interacting supersymmetric particles (sparticles) - the squarks and the gluinos [1]. However, a subset of these sparticles may still be rather light. An especially interesting possibility is the light stop scenario. In this case only one of the squarks - the lighter superpartner of the top quark or stop ($\widetilde{t}_1$) - is light and within the reach of the ongoing LHC experiments along with the electroweak sparticles (the sleptons and the electroweak gauginos)[3].

The light stop scenario is theoretically well-motivated. Due to a large mixing term - driven by the top quark mass - in the stop mass matrix, one of the mass eigenvalues (i.e $m_{\widetilde{t}_1}$) may turn out to be much smaller than the masses all other squarks. This result is model independent. Additional suppression of $m_{\widetilde{t}_1}$ may arise due to the underlying SUSY breaking mechanism. In the minimal supergravity (mSUGRA) model [8], e.g., all squarks have a common mass at the Grand Unification Theory (GUT) scale (M_G). Yet $m_{\widetilde{t}_1}$ at the weak scale may be suppressed relative to the masses of the other squarks due to renormalization group running between M_G and the weak scale driven by the large top quark Yukawa coupling. In this paper,however, we consider the light stop scenario at the weak scale as a phenomenological model without invoking any specific mechanism for the suppression of $m_{\widetilde{t}_1}$. Light stop squarks are cosmologically interesting since they can provide an explanation of baryogenesis [9]. This happens if $120 \lesssim m_{\widetilde{t}_1} \leq m_t$.

If $\widetilde{t}_1$ is the next to lightest super particle (NLSP) then another cosmologically interesting possibility opens up. The observed dark matter (DM) [10, 11] relic density of the universe [12] may be produced by the co-annihilation of $\widetilde{t}_1$ with the lightest supersymmetric particle (LSP), the DM candidate [13]. We shall consider models where the lightest neutralino ($\widetilde{\chi}_1^0$) is the LSP. For this co-annihilation, however, $\Delta m = m_{\widetilde{t}_1} - m_{\widetilde{\chi}_1^0}$ has to be rather small. The novel idea that the light stop scenario plays the major role in both baryogenesis and DM relic density production has also been emphasized in the literature [14]. In this work we shall focus on the possible constraints on the stop NLSP scenario for both large and small Δm from the published LHC 7-TeV data [2] corresponding to integrated luminosity $\mathcal{L} = 4.7$

---

[3] It has recently been emphasized [4, 5, 6] that if the squarks and the gluinos are compatible with the LHC mass bounds, the data is consistent with a wide variety of relatively light electroweak sparticles and they may have masses just above the corresponding lower limits from LEP [7].

fb^{-1}. Similar analyses were done for $\mathcal{L} = 1$ fb^{-1} data in [15]

Before starting the main discussion it is worthwhile to review the existing bounds on the stop NLSP from the LEP and the Tevatron. The Stop NLSP has the following decay modes: (i) $\widetilde{t}_1 \to c\widetilde{\chi}_1^0$ (the loop induced mode), (ii) $\widetilde{t}_1 \to bW\widetilde{\chi}_1^0$, (iii) $\widetilde{t}_1 \to t\widetilde{\chi}_1^0$ and (iv) $\widetilde{t}_1 \to bf\bar{f}'\widetilde{\chi}_1^0$

The Branching ratios of the modes (i) and (iv) are negligible if either (ii) or (iii) is kinematically accessible. The modes (i) and (iv) can in principle compete with one another [16], but the latter is negligible over most of the parameter space if tan β is moderately large (≥ 7) [16, 17] and will be ignored in this paper. If the $\widetilde{t}_1 - \widetilde{\chi}_1^0$ co-annihilation occurs then only decay mode (i) is allowed.

Most of the existing bounds on stop NLSP are derived by assuming that the loop induced decay (i) occur with 100%. The expected signal is (at least) 2 jets + $\not\!\!E_T$. For $m_{\widetilde{t}_1}$ within the kinematical reach of LEP, the bound is sensitive to small values of Δm as well and is given by $m_{\widetilde{t}_1} \geq 100$ (95.2) GeV for $\Delta m \geq 20$ (5) GeV for no mixing ($\theta_{\tilde{t}} = 0$) [7]. The numerical change in the limit is not drastic for other values of $\theta_{\tilde{t}}$. On the other hand the Tevatron experiments are sensitive to larger $m_{\widetilde{t}_1}$ but are valid for rather larger Δm. The best bound is $m_{\widetilde{t}_1} \geq 180$ GeV for $m_{\widetilde{\chi}_1^0} = 90$ GeV. Thus stop induced baryogenesis is disfavoured for Δm $\simeq 90$ GeV. For smaller Δm the efficiency of the kinematical selection requiring high p_T jets becomes rather poor due to the modest energy release in the decay process and the limit becomes weaker. The possibility of stop NLSP induced baryogenesis, therefore, remains open for small Δm. It is, also, fair to conclude that the possibility of $\widetilde{t}_1$ - LSP co-annihilation is only excluded for $m_{\widetilde{t}_1} \lesssim 100$ from the LEP bound.

The ATLAS or the CMS collaboration has not yet published any dedicated search for the pair production of the stop NLSP. However, this pair production followed by the decay of each stop in channels i) - iv) would certainly contribute to the jets + missing E_T signature carefully looked into by the LHC collaborations. The non-observation of this signal can potentially constrain the light stop scenario. It has, however, already been noted that the $\mathcal{L} \approx 1\text{fb}^{-1}$ data was sensitive to the stop sector [15]. However, we shall show below that the limits are much stronger for the $\mathcal{L} = 4.7$ fb^{-1} data. This large $\mathcal{L}$ accumulated at the LHC enables one to constrain the stop NLSP scenario meaningfully for smaller Δm and directly address the issue of Stop NLSP - LSP co-annihilation as a DM producing mechanism at a hadron collider.

We have computed the sparticle spectra and decay branching ratios (BRs) in the minimal supersymmetric extension of the standard model (MSSM) without any assumption regarding the soft breaking parameters using SUSPECT (v2.41) [18] and SDECAY [19]. Using PYTHIA (v6.424)[20] Monte Carlo (MC) event generator we have generated the jets + $\not{E}_T$ signal stemming from $\widetilde{t}_1\widetilde{t}_1^*$ events only. The next to leading order (NLO) cross-section for the $\widetilde{t}_1\widetilde{t}_1^*$ pair production have been computed by PROSPINO 2.1 [21] with CTEQ6.6M PDF [22]. The normalization and factorization scale is chosen to be $m_{\widetilde{t}_1}$. We have taken the pole mass of the top quark (running bottom quark mass evaluated in the $\overline{MS}$ scheme) as m_t (m_b) = 173.2 (4.25). We have computed the DM relic density using micrOMEGAs (v.2.4.1) [23]. The observed DM relic density (Ωh^2) in the universe measured by the WMAP collaboration [12] is given by $\Omega h^2 = 0.1126 \pm 0.0036$. If 10% theoretical uncertainty is added [24] then the DM relic density is bounded by $0.09 \leq \Omega h^2 \leq 0.13$ at 2σ level.



Figure 1: The distribution of N_{jet} (in the left panel) and p_T (in the right panel) of the hardest jet for the Benchmark points BP1 and BP2 (see text for details).

ATLAS Collaboration has updated their results for SUSY searches in $jets + \not{E}_T + 0l$ final state for $\mathcal{L} = 4.7$ fb^{-1} [2]. They have defined five inclusive analysis channels labelled as A to E according to jet multiplicity ($N_j = 2$ to 6). Further, the $N_j = 2$ sample is divided into two channels - A and A'. The channels and the details of kinemetical cuts for each of them are given in Table 1 of [2]. Depending upon the final cuts on the observable m_{eff}(incl.) (defined as the scalar sum of all jets with $P_T > 40$ GeV and $\not{E}_T$) each channel is further classified as

‘Tight’,‘Medium’ and ‘Loose’. In this way they have finally presented the results for 11 signal regions and have constrained any new physics model in terms of upper limit on the number of events N_{BSM} or effective cross-section σ_{BSM}/fb. The observed upper limits on N_{BSM} at 95 % Confidence Level (CL) for signal regions SRA-Tight, SRA-Medium, SRA’-Medium, SRB-Tight, SRC-Tight, SRC-Medium, SRC-Loose, SRD-Tight, SRE-Tight, SRE-Medium, SRE-Loose are 2.9, 25, 29, 3.1, 16, 18, 58, 10, 12, 12, 84 respectively [2].

We have adopted the selection criteria for different signal regions used by the ATLAS Collaboration [2]. Then we have checked whether the number of events from the $\widetilde{t}_1\widetilde{t}_1^*$ pair production followed by the decays (i) - (iii) of both the squarks exceed the corresponding upper bound for at least one of the signal regions to obtain the exclusion contour in the $m_{\widetilde{t}_1}$ - $m_{\widetilde{\chi}_1^0}$ plane.

If the BR of the decay mode $\widetilde{t}_1 \to c\widetilde{\chi}_1^0$ is 100 % then the signal will be 2j + $\not{E}_T$. In this case SRA-Medium or SRA’-Medium, specially designed for compressed spectra [2] is very effective in excluding the regions of the $m_{\widetilde{t}_1} - m_{\widetilde{\chi}_1^0}$ space. For small Δm, SRA’-Medium is the most sensitive one. On the otherhand, high jet multiplicity regions chosen by ATLAS are optimised for long decay chains [2]. For $\widetilde{t}_1 \to bW\widetilde{\chi}_1^0$ or $t\widetilde{\chi}_1^0$ decay, events will contain at least 2, 4 or 6 jets. Due to the lepton veto 2j signal will not be sensitive at all. In our analysis we have also checked that when BR of $\widetilde{t}_1 \to t\widetilde{\chi}_1^0$ is 100 % then the exclusion mainly comes from SRE-Medium ($N_j = 6$) channel. Again for $\widetilde{t}_1 \to bW\widetilde{\chi}_1^0$ effective signal regions are either SRC-Medium or SRE-Medium. The distributions of the number of jets (N_j) and the p_T of the hardest jet in the signal are shown in Fig. 1 for the two representative point BP1 and BP2. For both of them $m_{\widetilde{t}_1}$ is 200 GeV. But $m_{\widetilde{\chi}_1^0} = 180$ and 100 for BP1 and BP2 respectively. Both the distributions are drawn with nominal cuts and normalized to unity.

Fig. 2 depicts the regions in the $m_{\widetilde{t}_1}$ - $m_{\widetilde{\chi}_1^0}$ plane excluded by the ATLAS jets + $\not{E}_T$ data. In the region between the $m_{\widetilde{t}_1} = m_c + m_{\widetilde{\chi}_1^0}$ and the $m_{\widetilde{t}_1} = m_b + m_W + m_{\widetilde{\chi}_1^0}$ lines (Region-I), the loop decay (i) dominates. We have restricted ourselves to $(m_{\widetilde{t}_1})_{min} > m_c + m_{\widetilde{\chi}_1^0} + 2$ GeV. The same increment of 2 GeV has been introduced at each point representing a cross over from one region to another. In Region-II bounded by the lines $m_{\widetilde{t}_1} = m_b + m_W + m_{\widetilde{\chi}_1^0}$ and $m_{\widetilde{t}_1} = m_t + m_{\widetilde{\chi}_1^0}$ lines, the three body decay channel (ii) dominates. In regions I and II the parameter space to the left of the curved line is excluded. In Region-III below the line $m_{\widetilde{t}_1} = m_t + m_{\widetilde{\chi}_1^0}$, the two body decay mode (iii) is the only significant decay mode. The parameter space bounded by the dashed line and the x-axis is excluded. The following



Figure 2: Exclusion limits in the $m_{\widetilde{t}_1}$ - $m_{\widetilde{\chi}_1^0}$ plane for different decay modes of the $\widetilde{t}_1$ - NLSP. The x marked points represent the parameter space where $\widetilde{t}_1 - \widetilde{\chi}_1^0$ co-annihilation is viable.

points may be noted :

a) For very small Δm the jets in the signal essentially come from initial state and final state radiation. This result is, therefore , crucially dependent on the parton showering model in Pythia. In contrast for $\Delta m =$ 25 - 30 GeV, which is the case in the co-annihilation region, even the parton level p_T distribution develops a long tail. Our results for this region is seem to be more reliable. In the co-annihilation region $m_{\widetilde{t}_1} \lesssim 240$ GeV is disfavoured. In Region-II the best limit is $m_{\widetilde{t}_1} \gtrsim 230$ GeV for $m_{\widetilde{\chi}_1^0} \simeq 150$ GeV.

b) The $\widetilde{t}_1$ NLSP induced baryogenesis requiring $120 \lesssim m_{\widetilde{t}_1} \leq m_t$ seems to be disfavoured in the co-annihilation region. In Region-II $m_{\widetilde{t}_1} \lesssim 200$ GeV and hence, $\widetilde{t}_1$ induced baryogenesis is also disfavoured. In Region-III it is forbidden by definition.

c) Significant constraints are obtained in decay channels (ii) and (iii). The strong constraints obtained even for small Δm are due to the large $\mathcal{L}$ accumulated at the current LHC 7 TeV experiments. Most of the above bounds come from ATLAS selection criteria requiring two high p_T jets in the final state . That is why the constraints in Region-III are rather modest. Here the final state contains either large number of jets and/or at least one high p_T lepton which is vetoed.

References

[1] For reviews on Supersymmetry, see, *e.g.* , H. P. Nilles, Phys. Rep. **110**, 1 (1984); H. E. Haber and G. Kane, Phys. Rep. **117**, 75 (1985) ; J. Wess and J. Bagger, *Supersymmetry and Supergravity*, 2nd ed., (Princeton University Press, Princeton, 1991); M. Drees, P. Roy and R. M. Godbole, *Theory and Phenomenology of Sparticles*, (World Scientific, Singapore, 2005).

[2] ATLAS collaboration, ATLAS-CONF-2012-033

[3] CMS Collaboration, Report No. CMS-PAS-SUS-11-016.

[4] N. Bhattacharyya, A. Choudhury and A. Datta, Phys. Rev. D **83**, 115025 (2011).

[5] N. Bhattacharyya, A. Choudhury and A. Datta, Phys. Rev. D **84**, 095006 (2011).

[6] Arghya Choudhury and Amitava Datta, J. High Energy Phys. **06**, 2012 (006)

[7] For the latest limits on the sparticle masses from LEP experiments: see,e.g., http://lepsusy.web.cern.ch/lepsusy/.

[8] A. H. Chamseddine, R. Arnowitt and P. Nath, Phys. Rev. Lett. **49**, 970 (1982); R. Barbieri, S. Ferrara and C. A. Savoy, Phys. Lett. B **119**, 343 (1982); L. J. Hall, J. Lykken and S. Weinberg, Phys. Rev. D **27**, 2359 (1983); P. Nath, R. Arnowitt and A. H. Chamseddine, Nucl. Phys. B **227**, 121 (1983); N. Ohta, Prog. Theor. Phys. **70**, 542 (1983).

[9] M. Carena, M. Quiros and C. E. Wagner, Phys. Lett. B **380**, 81 (1996); D. Delepine, J. M. Gerard, R. Gonzalez Felipe and J. Weyers, Phys. Lett. B **386**, 183 (1996); J.

McDonald, Phys. Lett. B **413**, 30 (1997); J. M. Cline and G. D. Moore, Phys. Rev. Lett. **81**, 3315 (1998); M. Carena, G. Nardini, M. Quiros and C. E.M. Wagner, Nucl. Phys. B **812**, 243 (2009).

[10] W. L. Freedman and M. S. Turner, Rev. Mod. Phys. **75**, 1433 (2003); L. Roszkowski, Pramana **62**, 389 (2004). G. Bertone, D. Hooper and J. Silk, Phys. Rept. **405**, 279 (2005);

[11] H. Baer and X. Tata in *Physics at the Large Hadron Collider*, Indian National Science Academy, A Platinum Jubilee Special Issue (Eds. Amitava Datta, B. Mukhopadhyaya and A. Raychaudhuri; Springer, 2009).

[12] E. Komatsu *et al*, (WMAP Collaboration), Astrophys. J. Suppl. **192**, 18 (2011).

[13] C. Boehm, A. Djouadi and M. Drees, Phys. Rev. D **62**, 035012 (2000); J. R. Ellis, K. A. Olive and Y. Santoso, Astropart. Phys. 18, 395 (2003).

[14] C. Balazas, M. Carena and C. E. M. Wagner, Phys. Rev. D **70**, 015007 (2004); V. Cirigliano, S. Profamo, M. J. Ramsey Musolf J. High Energy Phys. **0607**, 002 (2006).

[15] N. Desai and B. Mukhopadhyaya, JHEP 05 (2012) 057; M. A. Ajaib, T. Li, Q. Shafi, Phys. Rev. D **85**, 055021 (2012); B. He, T. Li and Q. Shafi, JHEP 05 (2012) 148.

[16] C. Boehm, A. Djouadi and Y. Mambrini, Phys. Rev. D **61**, 095006 (2000).

[17] S.P. Das, Amitava Datta and M. Guchait, Phys. Rev. D **65**, 095006 (2002).

[18] A. Djouadi, J. L. Kneur and G. Moultaka, Comp. Phys. Comm. **176**, 426 (2007).

[19] M. Muhlleitner, A. Djouadi and Y. Mambrini, Comp. Phys. Comm. **168**, 46 (2005).

[20] T. Sjostrand, S. Mrenna and P. Skands, J. High Energy Phys. **0605**, 026 (2006).

[21] W. Beenakker *et al.*, Nucl. Phys. B **515**, 3 (1998).

[22] P. M. Nadolsky *et al*, Phys. Rev. D **78**, 013004 (2008).

[23] G. Belanger *et al*, Comp. Phys. Comm. **182**, 842 (2011).

[24] N. Baro, F. Boudjema and A. Semenov, Phys. Lett. B **660**, 550 (2008).